

q^2 interval [GeV ²]	$d\mathcal{B}(\text{B}_s^0 \rightarrow \phi\mu^+\mu^-)/\mathcal{B}(\text{B}_s^0 \rightarrow \text{J}/\psi\phi) dq^2$ [10 ⁻⁵ GeV ⁻²]	$d\mathcal{B}(\text{B}_s^0 \rightarrow \phi\mu^+\mu^-)/dq^2$ [10 ⁻⁸ GeV ⁻²]
1.1–2.5	2.51 ± 0.26 ± 0.07	2.58 ± 0.27 ± 0.07 ± 0.10
2.5–4	2.12 ± 0.23 ± 0.03	2.18 ± 0.23 ± 0.03 ± 0.08
4–6	2.48 ± 0.21 ± 0.03	2.55 ± 0.22 ± 0.03 ± 0.10
6–8	3.24 ± 0.24 ± 0.04	3.33 ± 0.25 ± 0.04 ± 0.13
11–12.5	4.78 ± 0.28 ± 0.08	4.93 ± 0.29 ± 0.08 ± 0.19
15–16	4.63 ± 0.33 ± 0.07	4.77 ± 0.34 ± 0.07 ± 0.18
16–19	4.14 ± 0.31 ± 0.18	4.26 ± 0.32 ± 0.18 ± 0.16
1.1–6	2.37 ± 0.13 ± 0.05	2.44 ± 0.14 ± 0.05 ± 0.09