

Search region	E_T^{miss} [GeV]	Lost lepton	$Z(\nu\nu)$	Rare	QCD	total SM	N_{data}
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) < 175 \text{ GeV}, N_j \geq 7, N_{\text{res}} \geq 1$							
10	250–400	420 ± 64	6.9 ± 2.1	12 ± 3	4.1 ± 2.7	443 ± 66	420
11	> 400	27 ± 5	1.6 ± 0.6	1.9 ± 0.6	0.15 ± 0.11	31 ± 6	28
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_j \geq 7, N_t = 0, N_{\text{res}} = 0, N_W = 0$							
12	250–450	148 ± 17	60 ± 8	14 ± 4	17 ± 9	239 ± 24	239
13	> 450	12 ± 3	13 ± 2	2.9 ± 0.8	1.2 ± 0.8	29 ± 4	35
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 1, N_{\text{res}} = 0, N_W = 0$							
14	> 650	$0.38^{+0.27}_{-0.22}$	0.71 ± 0.23	0.56 ± 0.17	0.03 ± 0.03	1.7 ± 0.4	4
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 0, N_{\text{res}} = 1, N_W = 0$							
15	250–450	170 ± 30	31 ± 8	23 ± 6	7.5 ± 5.1	231 ± 38	213
16	450–650	5.2 ± 1.5	5.0 ± 1.3	3.0 ± 0.8	0.58 ± 0.48	14 ± 3	8
17	> 650	$0.82^{+0.45}_{-0.34}$	0.8 ± 0.25	0.75 ± 0.35	$0.04^{+0.05}_{-0.04}$	$2.4^{+0.7}_{-0.6}$	2
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 0, N_{\text{res}} = 0, N_W = 1$							
18	350–550	13 ± 3	11 ± 3	3.6 ± 1.0	2.7 ± 1.8	30 ± 6	40
19	> 550	1.1 ± 0.4	1.9 ± 0.7	0.81 ± 0.42	0.03 ± 0.02	3.8 ± 1.0	2
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 1, N_{\text{res}} = 0, N_W = 1$							
20	> 550	0.21 ± 0.14	0.08 ± 0.05	0.1 ± 0.03	<0.01	0.38 ± 0.17	1
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 0, N_{\text{res}} = 1, N_W = 1$							
21	> 450	0.31 ± 0.19	0.21 ± 0.11	0.32 ± 0.1	<0.01	0.83 ± 0.3	0
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 1, N_{\text{res}} = 1, N_W = 0$							
22	> 450	0.01 ± 0.01	0.06 ± 0.04	0.2 ± 0.08	<0.01	0.28 ± 0.09	0
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t \geq 2, N_{\text{res}} = 0, N_W = 0$							
23	> 250	0.06 ± 0.06	<0.01	0.16 ± 0.07	<0.01	0.22 ± 0.1	1
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 0, N_{\text{res}} \geq 2, N_W = 0$							
24	> 250	1.9 ± 0.8	0.35 ± 0.22	1.5 ± 0.7	<0.01	3.8 ± 1.4	3
$N_b \geq 2, m_T(b_{1,2}, E_T^{\text{miss}}) > 175 \text{ GeV}, N_t = 0, N_{\text{res}} = 0, N_W \geq 2$							
25	> 250	1.5 ± 0.7	0.39 ± 0.2	0.17 ± 0.13	<0.01	2.1 ± 0.9	3