	$\widetilde{\chi}_1^{\pm} o \widetilde{\ell} \nu(\ell \widetilde{\nu}) o \ell \nu \widetilde{\chi}_1^0$	
Source of uncertainty	$(m_{\tilde{\chi}_1^{\pm}} = 800 \text{GeV}, m_{\tilde{\chi}_1^0} = 200 \text{GeV})$	
	Yields	$m_{\mathrm{T2}}(\ell\ell)$ shape
Integrated luminosity	1–3%	_
Trigger efficiency	2%	< 1%
Pileup	$\leq 4\%$	$\leq 4\%$
Jet energy scale	$\leq 5\%$	≤ 9%
Jet energy resolution	$\leq 3\%$	$\leq 14\%$
Unclustered energy	≤ 3%	$\leq 10\%$
Lepton ident./isolation	2–7%	≤ 3%
b tagging	< 1%	< 1%
b tagging (light jets)	< 1%	< 1%
Simulated samples statistics	3–15%	6–22%
Renorm./fact. scales	$\leq 1\%$	< 1%
Nonprompt leptons	< 1%	< 1%
Lepton ident./isolation (FASTSIM)	4.0%	
b tagging (FASTSIM)	< 1%	< 1%
ISR reweighting	$\leq 2\%$	$\leq 8\%$
$p_{\mathrm{T}}^{\mathrm{miss}}$ (FASTSIM)	≤ 11%	≤ 24%