Wilson coefficient	Best fit $[\text{TeV}^{-2}]$	68% CL [TeV ⁻²]	95% CL [TeV ⁻²]
$\bar{c}_{\mathrm{uW}}/\Lambda^2$	1.7	[-2.4, -0.5] and $[0.4, 2.4]$	[-2.9, 2.9]
$ \bar{c}_{\rm H}/\Lambda^2 - 16.8~{ m TeV^{-2}} $	15.6	[0,23.0]	[0,28.5]
$ \widetilde{c}_{3\mathrm{G}}/\Lambda^2 $	0.5	[0, 0.7]	[0, 0.9]
$\bar{c}_{3\mathrm{G}}/\Lambda^2$	-0.4	[-0.6, 0.1] and $[0.4, 0.7]$	[-0.7, 1.0]
$\bar{c}_{\mathrm{uG}}/\Lambda^2$	0.2	[0, 0.3]	[-1.0, -0.9] and $[-0.3, 0.4]$
$ \bar{c}_{\mathrm{uB}}/\Lambda^2 $	1.6	[0, 2.2]	[0, 2.7]
$\bar{c}_{\mathrm{Hu}}/\Lambda^2$	-9.3	[-10.3, -8.0] and $[0, 2.1]$	[-11.1, -6.5] and $[-1.6, 3.0]$
$\bar{c}_{2\mathrm{G}}/\Lambda^2$	0.4	[-0.9, -0.3] and $[-0.1, 0.6]$	[-1.1, 0.8]